Statistics

62Xxx

- [1] Joel E. Atkins and Gary J. Sherman, Sets of typical subsamples, Statist. Probab. Lett. 14 (1992), no. 2, 115–117. MR MR1173408 (93e:62008)
- [2] R. A. Bailey, Strata for randomized experiments, J. Roy. Statist. Soc. Ser. B **53** (1991), no. 1, 27–78, With discussion and a reply by the author. MR MR1094275 (92k:62134)
- [3] H. Evangelaras, I. Kotsireas, and C. Koukouvinos, Application of Gröbner bases to the analysis of certain two or three level factorial designs, Adv. Appl. Stat. 3 (2003), no. 1, 1–13. MR MR1983030 (2004d:62268)
- [4] H. Evangelaras and C. Koukouvinos, Combined arrays with minimum number of runs and maximum estimation efficiency, Comm. Statist. Theory Methods 33 (2004), no. 7, 1621–1628. MR MR2066372
- [5] Shuhong Gao, Guangran Jiang, and Mingfu Zhu, Solving the 100 Swiss Francs problem, 2008.
- [6] S. Georgiou and C. Koukouvinos, Some results on orthogonal designs and Hadamard matrices, Int. J. Appl. Math. 17 (2005), no. 4, 433–443. MR MR2199768 (2007b:05032)
- [7] S. Georgiou, C. Koukouvinos, and S. Stylianou, Construction of new skew Hadamard matrices and their use in screening experiments, Comput. Statist. Data Anal. 45 (2004), no. 3, 423–429. MR MR2050247
- [8] Alexey Koloydenko, Symmetric measures via moments, Bernoulli 14 (2008), no. 2, 362–390.
- [9] E. M. Rains, N. J. A. Sloane, and John Stufken, The lattice of N-run orthogonal arrays, J. Statist. Plann. Inference 102 (2002), no. 2, 477–500, Silver jubilee issue. MR MR1897221 (2003c:62138)